

2023; 16: e857

Carta al Editor

Polypharmacy in the elderly

Polimedicación en el adulto mayor

Dear Editor:

Scientific and technical development and the globalization of society have resulted in an increase in the number of available medicines and the need for information on their effectiveness and safety. Medicines are the most widely used medical technology in the contemporary world. (1)

Population aging is a situation facing humanity; it is associated with polypharmacy (condition defined by the World Health Organization as the use of five or more drugs simultaneously) ⁽²⁾ due to the multiple chronic diseases suffered by older adults.

Older adults are the population group that requires the most medication and requires the most medical attention 80 % of these patients have one or more non-communicable diseases and 36 % may suffer from more than three conditions that make it essential to use multiple medications, which leads to polypharmacy. This consumption leads to a high rate of adverse effects, drug interactions and mortality. (1)

In Spain, daily drug use ranges from two to three medications and more than 75 % consume them chronically. In Sweden, there is evidence of abuse of more than four different groups of drugs in adults over 85 years of age. In Mexico, the prevalence of polypharmacy in adults is 34 % and this percentage involves older adults. (1) In Cuba, 11 % of the elderly are on polypharmacy. (3)

The Cuban population is aging rapidly. This change in the age structure of a population has a direct impact on areas such as the economy, health and social security. ⁽³⁾ For this reason, the Family Doctor and Nurse Program (approved in 2023) points out as a fundamental activity the

http://revcimeq.sld.cu/index.php/imq revinmedquir@infomed.sld.cu

Under Creative Commons license







2023: 16: e857

achievement of healthy aging from a social, functional and psychological point of view, with the participation of the family and community factors. Another activity pointed out in this document is the prevention of damage to health caused by the use of multiple medications.

It is necessary for older adults to be advised by nursing professionals on the proper use of drugs, as well as to establish a work dynamic that must be addressed from an interdisciplinary, transdisciplinary and intersectoral perspective to create support systems for the proper use of medications.

Mary of Charity Ayala Viamontes ¹ https://orcid.org/0000/0002/1462/6313

Esther Izquierdo Mach ń ¹ https://orcid.org/0000/0002/7889/1814

Rolando Alvarez Marquez ² https://orcid.org/0000/0002/1650/3521

¹ "Lidia Doce" Nursing School. Havana, Cuba.
 ² Julio Trigo Lopez School of Medical Sciences. Havana, Cuba.

*Correspondence. Email: <u>eizquierdo@infomed.sld.cu</u>

BIBLIOGRAPHIC REFERENCES

1. Arroyo Gámez F, Heyerdahl Viau I, Pérez Sánchez R, Prado Galbarro F, Sánchez Herrera K, Mart nez Núñez J. Potenciales interacciones medicamentosas en un centro de salud público de México. Revista Cubana de Farmacia [Internet]. 2023 [citado 26 Jun 2024]; 56 (2): e892. Disponible en: https://revfarmacia.sld.cu/index.php/far/article/view/892

http://revcimeq.sld.cu/index.php/imq revinmedquir@infomed.sld.cu

Under Creative Commons license







2023; 16: e857

- 2. Triay-Diaz D, Bermúdez-Dom nguez M, Risco-Almenares M, León-Montero D. Vigilancia poscomercialización del VIDATOX® 30CH, Cuba 2013-2018. Revista Cubana de Farmacia [Internet]. 2020 [citado 15 Oct 2023]; 53 (2): e409 Disponible en: https://revfarmacia.sld.cu/index.php/far/article/view/409
- 3. Dáz-Molina M, Herrera-Preval Y. Prescripciones potencialmente inadecuadas en adultos mayores del municipio San Miguel del Padrón, La Habana, Cuba. Revista Cubana de Farmacia [Internet]. 2021 [citado 15 Oct 2023]; 54 (1): e508 Disponible en: https://revfarmacia.sld.cu/index.php/far/article/view/508

http://revcimeq.sld.cu/index.php/imq revinmedquir@infomed.sld.cu

Under Creative Commons license

